

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
29 March 2001 (29.03.2001)

PCT

(10) International Publication Number
WO 01/21609 A1(51) International Patent Classification⁷: C07D 311/22,
405/12, A61K 31/353, A61P 9/06

(21) International Application Number: PCT/JP00/06497

(22) International Filing Date:
22 September 2000 (22.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
11/269899 24 September 1999 (24.09.1999) JP(71) Applicant (for all designated States except US): NISSAN
CHEMICAL INDUSTRIES, LTD. [JP/JP]; 7-1, Kandan-
ishiki-cho 3-chome, Chiyoda-ku, Tokyo 101-0054 (JP).

(72) Inventors; and



(75) Inventors/Applicants (for US only): TANIKAWA,
Keizo [JP/JP]; Nissan Chemical Industries, Ltd., Central
Research Institute, 722-1, Tsuboi-cho, Funabashi-shi,
Chiba 274-8507 (JP). OHRAI, Kazuhiko [JP/JP]; Nissan
Chemical Industries, Ltd., Central Research Institute,
722-1, Tsuboi-cho, Funabashi-shi, Chiba 274-8507(JP). YANAGIHARA, Kazufumi [JP/JP]; Nissan
Chemical Industries, Ltd., Central Research Institute,
722-1, Tsuboi-cho, Funabashi-shi, Chiba 274-8507 (JP).
SHIGETA, Yukihiro [JP/JP]; Nissan Chemical Indus-
tries, Ltd., Central Research Institute, 722-1, Tsuboi-cho,
Funabashi-shi, Chiba 274-8507 (JP). TSUKAGOSHI,
Toru [JP/JP]; Nissan Chemical Industries, Ltd., Central
Research Institute, 722-1, Tsuboi-cho, Funabashi-shi,
Chiba 274-8507 (JP). YAMASHITA, Toru [JP/JP];
Nissan Chemical Industries, Ltd., Research Station of Bi-
ological Science, 1470, Ohaza Shiraoka, Shiraoka-machi,
Minamisaitama-gun, Saitama 349-0294 (JP).(74) Agents: HANABUSA, Tsuneo et al.; Hanabusa Patent Of-
fice, Ochanomizu Square B, 6, Kandasurugadai 1-chome,
Chiyoda-ku, Tokyo 101-0062 (JP).(81) Designated States (national): AU, CA, CN, CZ, HU, IL,
KR, LT, NO, NZ, RO, RU, SI, SK, UA, US.(84) Designated States (regional): European patent (AT, BE,
CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL,
PT, SE).

Published:

— With international search report.

[Continued on next page]

(54) Title: 4-OXYBENZOPYRAN DERIVATIVE



(57) Abstract: This invention relates to a 4-oxybenzopyran derivative of formula (I) wherein, R¹ and R² represent each independently a hydrogen atom, a C₁₋₆ alkyl group or a phenyl group; R³ represents a hydroxyl group or a C₁₋₆ alkylcarbonyloxy group; R⁴ represents a hydrogen atom, a C₃₋₆ cycloalkyl group, a C₁₋₆ alkyl group, a C₁₋₆ alkylcarbonyl group, a C₁₋₆ alkylaminocarbonyl group, a di-C₁₋₆ alkylaminocarbonyl group, an aryl group or a heteroaryl group; n represents an integer of 0-4; X represents -C(=O)NR⁷-, -NR⁸-, -NHC(=O)NH- or -S(O)₂NH-; R⁵ represents a hydrogen atom or (a), (b), (c), (d), (e), (f), (g), (h), (i), (j), (k), or (l); R⁶ represents a hydrogen atom, a halogen atom, a nitro group or a cyano group; or a pharmaceutically acceptable salt thereof. And this invention also relates to an antiarrhythmic agent having the prolongation effect on the functional refractory period comprising said compound or a pharmaceutically acceptable salt thereof as an active ingredient.



— *Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

DESCRIPTION

4-OXYBENZOPYRAN DERIVATIVE

Technical Field

The present invention relates to 4-oxybenzopyran derivatives having a prolongation effect on the functional refractory period, which are used for treatments of arrhythmia in mammal including human beings.

Background Art

As benzopyran derivatives, there have been known 4-acylaminobenzopyran derivatives exemplified by Cromakalim (Japanese Patent Application Laid-Open No. Sho 58-67683). These 4-acylaminobenzopyran derivatives exemplified by Cromakalim are known to open an ATP sensitive K^+ channel and to be effective for the treatment of hypertension or asthma, but there has not been any mention as to the treatment for arrhythmia based on the prolongation effect on the functional refractory period.

Now, conventional antiarrhythmic agents having the prolongation effect on the functional refractory period as a main function (such as Class I drugs of antiarrhythmic agent classification according to Vaughan Williams, or d-sotalol belonging to Class III) have highly dangerous arrhythmic inducing actions that can result in sudden death such as torsades de pointes based on extension of ventricular muscle action potential relating to the prolongation effect on the functional refractory period, which become the therapeutic problems. Thus, agents having less side effects are desired.


The inventors of the present invention have made an intensive study of compounds having the prolongation effect on the functional refractory period more selective for atrium muscle than for ventricular muscle, and found that the compound of the general formula (I) has a prolongation effect on the functional refractory period selective for atrium muscle without any influence on the refractory period of ventricular muscle and action potential

parameters.

Disclosure of Invention

The inventors of the present invention have studied eagerly 4-oxybenzopyran derivatives, and found that the compound of the formula (I) has the strong prolongation effect on the functional refractory period, and it is useful as an antiarrhythmic agent. The present invention has been made based on this finding.

The present invention relates to a 4-oxybenzopyran derivative of the formula (I)



wherein, R^1 and R^2 represent each independently a hydrogen atom; a C_{1-6} alkyl group in which said alkyl group may be optionally substituted with a halogen atom, a C_{1-6} alkoxy group or a hydroxyl group; or a phenyl group in which said phenyl group may be optionally substituted with a halogen atom, a hydroxyl group, a nitro group, a cyano group, a C_{1-6} alkyl group or a C_{1-6} alkoxy group;

R^3 represents a hydroxyl group or a C_{1-6} alkylcarbonyloxy group;


R^4 represents a hydrogen atom, a C_{3-6} cycloalkyl group, a C_{1-6} alkyl group, a C_{1-6} alkylcarbonyl group, a C_{1-6} alkylaminocarbonyl group or a di- C_{1-6} alkylaminocarbonyl group in which said C_{1-6} alkyl group, said C_{1-6} alkylcarbonyl group, said C_{1-6} alkylaminocarbonyl group and said di- C_{1-6} alkylaminocarbonyl group may be each optionally substituted with a halogen atom, a C_{1-6} alkoxy group, a C_{1-6} alkoxy group substituted by a halogen atom; a carboxyl group, a C_{1-6} alkoxycarbonyl group, a hydroxyl group, or an aryl group or a heteroaryl group; in which said aryl group and said heteroaryl group may be optionally substituted with $(R^9)_x$, in which R^9 represents a halogen atom, a hydroxyl group, a C_{1-6} alkyl group, a C_{1-6} alkyl group substituted by a halogen atom or a C_{1-6} alkoxy

group; a C₁₋₆ alkoxy group, a C₁₋₆ alkoxy group substituted by a halogen atom; or R⁹ represents a nitro group, a cyano group, a formyl group, a formamide group, an amino group, a C₁₋₆ alkylamino group, a di-C₁₋₆ alkylamino group, a C₁₋₆ alkylcarbonylamino group, a C₁₋₆ alkylsulfonylamino group, an aminocarbonyl group, a C₁₋₆ alkylaminocarbonyl group, a di-C₁₋₆ alkylaminocarbonyl group, a C₁₋₆ alkylcarbonyl group, a C₁₋₆ alkoxycarbonyl group, an aminosulfonyl group, a C₁₋₆ alkylsulfonyl group, a carboxyl group or an arylcarbonyl group, r represents an integer of 1-3 and each R⁹ may same or different if r represents 2 or 3; or R⁴ represents an aryl group or a heteroaryl group in which said aryl group and said heteroaryl group may be optionally substituted with (R¹⁰)_q in which R¹⁰ has the same meaning as R⁹, q represents an integer of 1-3, and each R¹⁰ may be same or different if q represents 2 or 3;

n represents an integer of 0-4;

X represents -C(=O)NR⁷-, -NR⁸-, -NHC(=O)NH- or -S(O)₂NH- (in which R⁷ and R⁸ represent each independently a hydrogen atom or a C₁₋₆ alkyl group);

R⁵ represents a hydrogen atom; or a group of the formula:



in which R^{12} represents a hydrogen atom; a halogen atom; a C_{1-6} alkyl group in which the alkyl group may be optionally substituted with a halogen atom or a C_{1-6} alkoxy group; a C_{1-6} alkoxy group in which said alkoxy group may be optionally substituted with a halogen atom; a phenyl group in which said phenyl group may be optionally substituted with a halogen atom, a hydroxyl group, a C_{1-6} alkyl group or a C_{1-6} alkoxy group; or R^{12} represents a hydroxyl group, a nitro group, a cyano group, a formyl group, a formamide group, an amino group, a C_{1-6} alkylamino group, a di- C_{1-6} alkylamino group, a C_{1-6} alkylcarbonylamino group, a C_{1-6} alkylsulfonylamino group, an aminocarbonyl group, a C_{1-6} alkylaminocarbonyl group, a di- C_{1-6} alkylaminocarbonyl group, a C_{1-6} alkylcarbonyl group, a C_{1-6} alkoxycarbonyl group, an aminosulfonyl group, a C_{1-6} alkylsulfonyl group, a carboxyl group or an arylcarbonyl group, m represents an integer of 1-3, and each R^{12} may be same or different if m represents 2 or 3, and R^{13} represents a hydrogen atom or a C_{1-6} alkyl group; and

R^6 represents a hydrogen atom, a halogen atom, a nitro group or a cyano group;
or a pharmaceutically acceptable salt thereof.

The compound according to the present invention has the strong prolongation effect on the functional refractory period and it can be used as a drug for treating arrhythmia.

Respective substituents for the compound (I) according to the present invention are illustrated specifically as follows.

Herein, "n" means normal, "i" means iso, "s" means secondary, "t" means tertiary, "c" means cyclo, "o" means ortho, "m" means meta, and "p" means para.

As C_{1-6} alkyl groups, there may be mentioned methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, 1-pentyl, 2-pentyl, 3-pentyl, i-pentyl, neopentyl, 2,2-dimethylpropyl, 1-hexyl, 2-hexyl, 3-hexyl, 1-methyl-n-pentyl, 1,1,2-trimethyl-n-propyl, 1,2,2-trimethyl-n-propyl, 3,3-dimethyl-n-butyl, trifluoromethyl, trifluoroethyl, pentafluoroethyl, cyanomethyl and hydroxymethyl, etc.

Preferably, there may be mentioned methyl, ethyl, n-propyl,

i-propyl and n-butyl.

As halogen atoms, there may be mentioned a fluorine atom, a chlorine atom, a bromine atom and an iodine atom. Preferably, there may be mentioned a fluorine atom, a chlorine atom and a bromine atom.

As C₁₋₆ alkoxy groups, there may be mentioned methoxy, trifluoromethoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, i-butoxy, s-butoxy, t-butoxy, 1-pentyloxy, 2-pentyloxy, 3-pentyloxy, i-pentyloxy, neopentyloxy, 2,2-dimethylpropoxy, 1-hexyloxy, 2-hexyloxy, 3-hexyloxy, 1-methyl-n-pentyloxy, 1,1,2-trimethyl-n-propoxy, 1,2,2-trimethyl-n-propoxy and 3,3-dimethyl-n-butoxy, etc.

Preferably, there may be mentioned methoxy, ethoxy, n-propoxy and i-propoxy.

As C₁₋₆ alkylcarbonyloxy groups, there may be mentioned methylcarbonyloxy, ethylcarbonyloxy, n-propylcarbonyloxy, i-propylcarbonyloxy, n-butylcarbonyloxy, i-butylcarbonyloxy, s-butylcarbonyloxy, t-butylcarbonyloxy, 1-pentylcarbonyloxy, 2-pentylcarbonyloxy, 3-pentylcarbonyloxy, i-pentylcarbonyloxy, neopentylcarbonyloxy, t-pentylcarbonyloxy, 1-hexylcarbonyloxy, 2-hexylcarbonyloxy, 3-hexylcarbonyloxy, 1-methyl-n-pentylcarbonyloxy, 1,1,2-trimethyl-n-propylcarbonyloxy, 1,2,2-trimethyl-n-propylcarbonyloxy and 3,3-dimethyl-n-butylcarbonyloxy, etc.

Preferably, there may be mentioned methylcarbonyloxy, ethylcarbonyloxy, n-propylcarbonyloxy, i-propylcarbonyloxy, n-butylcarbonyloxy and t-butylcarbonyloxy.

As C₃₋₆ cycloalkyl groups, there may be mentioned cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl, etc.

Preferably, there may be mentioned cyclopropyl, cyclobutyl and cyclohexyl.

As C₁₋₆ alkylcarbonyl groups, there may be mentioned methylcarbonyl, ethylcarbonyl, n-propylcarbonyl, i-propylcarbonyl, n-butylcarbonyl, i-butylcarbonyl, s-butylcarbonyl, t-butylcarbonyl, 1-pentylcarbonyl, 2-pentylcarbonyl, 3-pentyl-

carbonyl, i-pentylcarbonyl, neopentylcarbonyl, t-pentylcarbonyl, 1-hexylcarbonyl, 2-hexylcarbonyl and 3-hexylcarbonyl.

Preferably, there may be mentioned methylcarbonyl, ethylcarbonyl, n-propylcarbonyl, i-propylcarbonyl and n-butylcarbonyl.

As C₁₋₆ alkylaminocarbonyl groups, there may be mentioned methylaminocarbonyl, ethylaminocarbonyl, n-propylaminocarbonyl, i-propylaminocarbonyl, n-butylaminocarbonyl, i-butylaminocarbonyl, s-butylaminocarbonyl, t-butylaminocarbonyl, 1-pentylaminocarbonyl, 2-pentylaminocarbonyl, 3-pentylaminocarbonyl, i-pentylaminocarbonyl, neopentylaminocarbonyl, t-pentylaminocarbonyl, 1-hexylaminocarbonyl, 2-hexylaminocarbonyl and 3-hexylaminocarbonyl, etc.

Preferably, there may be mentioned methylaminocarbonyl, ethylaminocarbonyl, n-propylaminocarbonyl, i-propylaminocarbonyl and n-butylaminocarbonyl.

As di-C₁₋₆ alkylaminocarbonyl groups, there may be mentioned dimethylaminocarbonyl, diethylaminocarbonyl, di-n-propylaminocarbonyl, di-i-propylaminocarbonyl, di-c-propylaminocarbonyl, di-n-butylaminocarbonyl, di-i-butylaminocarbonyl, di-s-butylaminocarbonyl, di-t-butylaminocarbonyl, di-c-butylaminocarbonyl, di-1-pentylaminocarbonyl, di-2-pentylaminocarbonyl, di-3-pentylaminocarbonyl, di-i-pentylaminocarbonyl, di-neopentylaminocarbonyl, di-t-pentylaminocarbonyl, di-c-pentylaminocarbonyl, di-1-hexylaminocarbonyl, di-2-hexylaminocarbonyl and di-3-hexylaminocarbonyl, etc.

Preferably, there may be mentioned dimethylaminocarbonyl, diethylaminocarbonyl, di-n-propylaminocarbonyl, di-i-propylaminocarbonyl, di-c-propylaminocarbonyl and di-n-butylaminocarbonyl.

As C₁₋₆ alkoxy carbonyl groups, there may be mentioned methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, i-propoxycarbonyl, n-butoxycarbonyl, i-butoxycarbonyl, s-butoxycarbonyl, t-butoxycarbonyl, 1-pentyloxycarbonyl, 2-pentyloxycarbonyl, 3-pentyloxycarbonyl, i-pentyloxycarbonyl, neopentyloxycarbonyl, t-pentyloxycarbonyl, 1-hexyloxycarbonyl,

2-hexyloxycarbonyl and 3-hexyloxycarbonyl, etc.

Preferably, there may be mentioned methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, i-propoxycarbonyl, n-butoxycarbonyl, i-butoxycarbonyl, s-butoxycarbonyl and t-butoxycarbonyl.

As aryl groups, there may be mentioned phenyl, biphenyl, naphthyl, anthryl and phenanthryl etc.

Preferably, there may be mentioned phenyl, biphenyl and naphthyl.

As heteroaryl groups, there may be mentioned 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, 2-pyranyl, 3-pyranyl, 4-pyranyl, 2-benzofuranyl, 3-benzofuranyl, 4-benzofuranyl, 5-benzofuranyl, 6-benzofuranyl, 7-benzofuranyl, 1-isobenzofuranyl, 4-isobenzofuranyl, 5-isobenzofuranyl, 2-benzothienyl, 3-benzothienyl, 4-benzothienyl, 5-benzothienyl, 6-benzothienyl, 7-benzothienyl, 1-isobenzothienyl, 4-isobenzothienyl, 5-isobenzothienyl, 2-chromenyl, 3-chromenyl, 4-chromenyl, 5-chromenyl, 6-chromenyl, 7-chromenyl, 8-chromenyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 1-imidazolyl, 2-imidazolyl, 4-imidazolyl, 1-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 3-pyridazinyl, 4-pyridazinyl, 1-indoliziny, 2-indoliziny, 3-indoliziny, 5-indoliziny, 6-indoliziny, 7-indoliziny, 8-indoliziny, 1-isoindolyl, 4-isoindolyl, 5-isoindolyl, 1-indolyl, 2-indolyl, 3-indolyl, 4-indolyl, 5-indolyl, 6-indolyl, 7-indolyl, 1-indazolyl, 2-indazolyl, 3-indazolyl, 4-indazolyl, 5-indazolyl, 6-indazolyl, 7-indazolyl, 1-puriny, 2-puriny, 3-puriny, 6-puriny, 7-puriny, 8-puriny, 2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl, 7-quinolyl, 8-quinolyl, 1-isoquinolyl, 3-isoquinolyl, 4-isoquinolyl, 5-isoquinolyl, 6-isoquinolyl, 7-isoquinolyl, 8-isoquinolyl, 1-phthalazinyl, 5-phthalazinyl, 6-phthalazinyl, 2-naphthyridinyl, 3-naphthyridinyl, 4-naphthyridinyl, 2-quinoxaliny, 5-quinoxaliny, 6-quinoxaliny,

2-quinazolinyl, 4-quinazolinyl, 5-quinazolinyl, 6-quinazolinyl, 7-quinazolinyl, 8-quinazolinyl, 3-cinnolinyl, 4-cinnolinyl, 5-cinnolinyl, 6-cinnolinyl, 7-cinnolinyl, 8-cinnolinyl, 2-pteridiny, 4-pteridiny, 6-pteridiny, 7-pteridiny and 3-furazany, etc.

Preferably, there may be mentioned 2-pyridyl, 3-pyridyl and 4-pyridyl, etc.

As C₁₋₆ alkylamino groups, there may be mentioned methylamino, ethylamino, n-propylamino, i-propylamino, c-propylamino, n-butylamino, i-butylamino, s-butylamino, t-butylamino, c-butylamino, 1-pentylamino, 2-pentylamino, 3-pentylamino, i-pentylamino, neopentylamino, t-pentylamino, c-pentylamino, 1-hexylamino, 2-hexylamino, 3-hexylamino, c-hexylamino, 1-methyl-n-pentylamino, 1,1,2-trimethyl-n-propylamino, 1,2,2-trimethyl-n-propylamino and 3,3-dimethyl-n-butylamino, etc.

Preferably, there may be mentioned methylamino, ethylamino, n-propylamino, i-propylamino and n-butylamino.

As di-C₁₋₆ alkylamino groups, there may be mentioned dimethylamino, diethylamino, di-n-propylamino, di-i-propylamino, di-c-propylamino, di-n-butylamino, di-i-butylamino, di-s-butylamino, di-t-butylamino, di-c-butylamino, di-1-pentylamino, di-2-pentylamino, di-3-pentylamino, di-i-pentylamino, di-neopentylamino, di-t-pentylamino, di-c-pentylamino, di-1-hexylamino, di-2-hexylamino, di-3-hexylamino, di-c-hexylamino, di-(1-methyl-n-pentyl)amino, di-(1,1,2-trimethyl-n-propyl)amino, di-(1,2,2-trimethyl-n-propyl)amino, di-(3,3-dimethyl-n-butyl)amino, methyl(ethyl)amino, methyl(n-propyl)amino, methyl(i-propyl)amino, methyl(c-propyl)amino, methyl(n-butyl)amino, methyl(i-butyl)amino, methyl(s-butyl)amino, methyl(t-butyl)amino, methyl(c-butyl)amino, ethyl(n-propyl)amino, ethyl(i-propyl)amino, ethyl(c-propyl)amino, ethyl(n-butyl)amino, ethyl(i-butyl)amino, ethyl(s-butyl)amino, ethyl(t-butyl)amino, ethyl(c-butyl)amino, n-propyl(i-propyl)amino, n-propyl(c-propyl)amino, n-propyl(n-butyl)amino, n-propyl(i-butyl)amino, n-propyl(s-butyl)amino, n-propyl(t-butyl)amino, n-propyl(c-butyl)amino, i-propyl(c-propyl)amino, i-propyl(n-butyl)amino,

i-propyl(i-butyl)amino, i-propyl(s-butyl)amino, i-propyl(t-butyl)amino, i-propyl(c-butyl)amino, c-propyl(n-butyl)amino, c-propyl(i-butyl)amino, c-propyl(s-butyl)amino, c-propyl(t-butyl)amino, c-propyl(c-butyl)amino, n-butyl(i-butyl)amino, n-butyl(s-butyl)amino, n-butyl(t-butyl)amino, n-butyl(c-butyl)amino, i-butyl(s-butyl)amino, i-butyl(t-butyl)amino, i-butyl(c-butyl)amino, s-butyl(t-butyl)amino, s-butyl(c-butyl)amino and t-butyl(c-butyl)amino, etc.

Preferably, there may be mentioned dimethylamino, diethylamino, di-n-propylamino, di-i-propylamino and di-n-butylamino.

As C₁₋₆ alkylcarbonylamino groups, there may be mentioned methylcarbonylamino, ethylcarbonylamino, n-propylcarbonylamino, i-propylcarbonylamino, n-butylcarbonylamino, i-butylcarbonylamino, s-butylcarbonylamino, t-butylcarbonylamino, 1-pentylcarbonylamino, 2-pentylcarbonylamino, 3-pentylcarbonylamino, i-pentylcarbonylamino, neopentylcarbonylamino, t-pentylcarbonylamino, 1-hexylcarbonylamino, 2-hexylcarbonylamino and 3-hexylcarbonylamino, etc.

Preferably, there may be mentioned methylcarbonylamino, ethylcarbonylamino, n-propylcarbonylamino, i-propylcarbonylamino and n-butylcarbonylamino.

As C₁₋₆ alkylsulfonylamino groups, there may be mentioned methylsulfonylamino, ethylsulfonylamino, n-propylsulfonylamino, i-propylsulfonylamino, n-butylsulfonylamino, i-butylsulfonylamino, s-butylsulfonylamino, t-butylsulfonylamino, 1-pentylsulfonylamino, 2-pentylsulfonylamino, 3-pentylsulfonylamino, i-pentylsulfonylamino, neopentylsulfonylamino, t-pentylsulfonylamino, 1-hexylsulfonylamino, 2-hexylsulfonylamino and 3-hexylsulfonylamino, etc.

Preferably, there may be mentioned methylsulfonylamino, ethylsulfonylamino, n-propylsulfonylamino, i-propylsulfonylamino and n-butylsulfonylamino.

As C₁₋₆ alkylsulfonyl groups, there may be mentioned methanesulfonyl and ethanesulfonyl.


As arylcarbonyl groups, there may be mentioned benzoyl,

p-methylbenzoyl, p-t-butylbenzoyl, p-methoxybenzoyl, p-chlorobenzoyl, p-nitrobenzoyl and p-cyanobenzoyl.

Preferably, there may be mentioned benzoyl, p-nitrobenzoyl and p-cyanobenzoyl.

As preferable compounds used in the present invention, the following compounds may be mentioned.


- (1) A 4-oxybenzopyran derivative of the formula (I) or pharmaceutically acceptable salt thereof, wherein R^1 and R^2 represent both methyl groups and R^3 represents a hydroxyl group.
- (2) A 4-oxybenzopyran derivative or pharmaceutically acceptable salt according to the aforementioned (1), wherein R^6 represents a hydrogen atom or a nitro group.
- (3) A 4-oxybenzopyran derivative or pharmaceutically acceptable salt thereof according to the aforementioned (2), wherein X represents $-C(=O)NH-$, $-C(=O)NMe-$, $-S(O)_2NH-$ or $-NHC(=O)NH-$.
- (4) A 4-oxybenzopyran derivative or pharmaceutically acceptable salt thereof according to the aforementioned (3), wherein X represents $-C(=O)NH-$ and R^5 represents



- (5) A 4-oxybenzopyran derivative or pharmaceutically acceptable salt thereof according to the aforementioned (4), wherein R^4 represents a C_{1-6} alkyl group.


Specific examples of the compounds that can be used in the present invention are shown as follows, but the present invention is not limited thereto. Herein, "Me" means a methyl group, "Et" means an ethyl group, "Pr" means a propyl group, "Bu" means a butyl group, "Pen" means a pentyl group, "Hex" means a hexyl group, "Ph" means a phenyl group, "Ac" means an acetyl group ($COCH_3$), and "-" means a bond, respectively.

Table 1




R ¹	R ²	R ³	R ⁴	R ⁶	n
Me	Me	OH	H	H	1
H	H	OH	c-Pr	H	1
Me	Me	OH	c-Hex	H	2
Me	Me	OH	Me	H	1
Me	Me	OH	Et	H	1
Me	Me	OH	n-Pr	H	1
Me	Me	OH	i-Pr	H	1
Me	Me	OH	n-Bu	H	2
Me	Me	OH	n-Pen	H	3
Me	Me	OH	n-Hex	H	4
Me	Me	OH	COMe	H	2
Me	Me	OH	CONHMe	H	2
Me	Me	OH	CONMe ₂	H	2
Me	Me	OCOMe	CF ₃	H	2
Me	Me	OCOEt	CH ₂ Ph	NO ₂	2
Me	Me	OH	CH ₂ CH ₂ Ph	NO ₂	1
Me	Me	OH	CH ₂ CH ₂ Ph	NO ₂	2
Me	Me	OH	CH ₂ CH ₂ Ph	NO ₂	3
Me	Me	OH	CH ₂ CH ₂ Ph	NO ₂	4
Et	Et	OH	CH ₂ CH ₂ Ph	NO ₂	2
n-Pr	n-Pr	OH	CH ₂ CH ₂ Ph	NO ₂	2
i-Pr	i-Pr	OH	CH ₂ CH ₂ Ph	NO ₂	2
n-Bu	n-Bu	OH	CH ₂ CH ₂ Ph	NO ₂	2
i-Bu	i-Bu	OH	CH ₂ CH ₂ Ph	NO ₂	2
t-Bu	t-Bu	OH	CH ₂ CH ₂ Ph	NO ₂	3
n-Pen	n-Pen	OH	CH ₂ CH ₂ Ph	NO ₂	3
n-Hex	n-Hex	OH	CH ₂ CH ₂ Ph	NO ₂	3
CF ₃	CF ₃	OH	CH ₂ CH ₂ Ph	NO ₂	3
CH ₂ OCH ₃	CH ₂ OCH ₃	OH	CH ₂ CH ₂ Ph	NO ₂	3

Table 2




R^1	R^2	R^3	R^4	R^6	n
H	H	OH	H	H	0
Me	Me	OH	c-Pr	H	1
Me	Me	OH	c-Hex	H	2
Me	Me	OH	Me	Cl	3
Me	Me	OH	Et	Br	4
Me	Me	OH	n-Pr	H	0
Me	Me	OH	i-Pr	H	1
Me	Me	OH	n-Bu	H	2
Me	Me	OH	n-Pen	H	3
Me	Me	OH	n-Hex	F	4
Me	Me	OH	COMe	NO_2	2
Me	Me	OH	CONHMe	NO_2	2
Me	Me	OH	CONMe ₂	NO_2	2
Me	Me	OCOMe	CF_3	H	2
Me	Me	OCOEt	CH_2Ph	CN	2
Me	Me	OH	$\text{CH}_2\text{CH}_2\text{Ph}$	CN	1
Me	Me	OH	$\text{CH}_2\text{CH}_2\text{Ph}$	CN	2
Me	Me	OH	$\text{CH}_2\text{CH}_2\text{Ph}$	CN	3
Me	Me	OH	$\text{CH}_2\text{CH}_2\text{Ph}$	CN	4
Et	Et	OH	$\text{CH}_2\text{CH}_2\text{Ph}$	CN	2
n-Pr	n-Pr	OH	$\text{CH}_2\text{CH}_2\text{Ph}$	CN	2
i-Pr	i-Pr	OH	$\text{CH}_2\text{CH}_2\text{Ph}$	CN	2
n-Bu	n-Bu	OH	Ph	CN	2
i-Bu	i-Bu	OH	Ph	F	2
t-Bu	t-Bu	OH	Ph	F	3
n-Pen	n-Pen	OH	Ph	Cl	3
n-Hex	n-Hex	OH	Ph	Cl	3
CF_3	CF_3	OH	Ph	Br	3
CH_2OCH_3	CH_2OCH_3	OH	Ph	Br	3

Table 3




R^1	R^2	R^3	R^4	R^6	n
H	H	OH	H	H	0
Me	Me	OH	c-Pr	H	1
Me	Me	OH	c-Hex	H	2
Me	Me	OH	Me	H	3
Me	Me	OH	Et	H	4
Me	Me	OH	n-Pr	H	0
Me	Me	OH	i-Pr	H	1
Me	Me	OH	n-Bu	H	2
Me	Me	OH	n-Pen	H	3
Me	Me	OH	n-Hex	H	4
Me	Me	OH	COMe	H	2
Me	Me	OH	CONHMe	H	2
Me	Me	OH	CONMe ₂	H	2
Me	Me	OCOMe	CF ₃	H	2
Me	Me	OCOEt	CH ₂ Ph	NO ₂	2
Me	Me	OH	CH ₂ CH ₂ Ph	NO ₂	1
Me	Me	OH	CH ₂ CH ₂ Ph	NO ₂	2
Me	Me	OH	CH ₂ CH ₂ Ph	NO ₂	3
Me	Me	OH	CH ₂ CH ₂ Ph	NO ₂	4
Et	Et	OH	CH ₂ CH ₂ Ph	NO ₂	2
n-Pr	n-Pr	OH	CH ₂ CH ₂ Ph	NO ₂	2
i-Pr	i-Pr	OH	CH ₂ CH ₂ Ph	NO ₂	2
n-Bu	n-Bu	OH	CH ₂ CH ₂ Ph	NO ₂	2
i-Bu	i-Bu	OH	CH ₂ CH ₂ Ph	NO ₂	2
t-Bu	t-Bu	OH	CH ₂ CH ₂ Ph	NO ₂	3
n-Pen	n-Pen	OH	CH ₂ CH ₂ Ph	NO ₂	3
n-Hex	n-Hex	OH	CH ₂ CH ₂ Ph	NO ₂	3
CF ₃	CF ₃	OH	CH ₂ CH ₂ Ph	NO ₂	3
CH ₂ OCH ₃	CH ₂ OCH ₃	OH	CH ₂ CH ₂ Ph	NO ₂	3

Table 4




R ¹	R ²	R ³	R ⁴	R ⁶	n
H	H	OH	H	H	0
Me	Me	OH	c-Pr	H	1
Me	Me	OH	c-Hex	H	2
Me	Me	OH	Me	H	3
Me	Me	OH	Et	H	4
Me	Me	OH	n-Pr	H	0
Me	Me	OH	i-Pr	H	1
Me	Me	OH	n-Bu	H	2
Me	Me	OH	n-Pen	H	3
Me	Me	OH	n-Hex	H	4
Me	Me	OH	COMe	H	2
Me	Me	OH	CONHMe	H	2
Me	Me	OH	CONMe ₂	H	2
Me	Me	OCOMe	CF ₃	H	2
Me	Me	OCOEt	CH ₂ Ph	NO ₂	2
Me	Me	OH	CH ₂ CH ₂ Ph	NO ₂	1
Me	Me	OH	CH ₂ CH ₂ Ph	NO ₂	2
Me	Me	OH	CH ₂ CH ₂ Ph	NO ₂	3
Me	Me	OH	CH ₂ CH ₂ Ph	NO ₂	4
Et	Et	OH	CH ₂ CH ₂ Ph	NO ₂	2
n-Pr	n-Pr	OH	CH ₂ CH ₂ Ph	NO ₂	2
i-Pr	i-Pr	OH	CH ₂ CH ₂ Ph	NO ₂	2
n-Bu	n-Bu	OH	CH ₂ CH ₂ Ph	NO ₂	2
i-Bu	i-Bu	OH	CH ₂ CH ₂ Ph	NO ₂	2
t-Bu	t-Bu	OH	CH ₂ CH ₂ Ph	NO ₂	3
n-Pen	n-Pen	OH	CH ₂ CH ₂ Ph	NO ₂	3
n-Hex	n-Hex	OH	CH ₂ CH ₂ Ph	NO ₂	3
CF ₃	CF ₃	OH	CH ₂ CH ₂ Ph	NO ₂	3
CH ₂ OCH ₃	CH ₂ OCH ₃	OH	CH ₂ CH ₂ Ph	NO ₂	3

Table 5




R ⁴	R ⁶	R ¹²	n
H	H	p-MeO	0
c-Pr	H	p-MeO	1
c-Hex	H	p-MeO	1
Me	Cl	p-MeO	1
Et	Br	p-MeO	1
n-Pr	H	m-MeO	1
i-Pr	H	o-MeO	1
n-Bu	H	p-Me	1
n-Pen	H	p-Et	1
n-Hex	F	m-Et	1
COMe	NO ₂	o-Et	1
CONHMe	NO ₂	p-Cl	2
CONMe ₂	NO ₂	p-F	2
CF ₃	H	p-Ph	3
CH ₂ Ph	CN	p-OH	3
CH ₂ CH ₂ Ph	CN	p-NO ₂	4
CH ₂ CH ₂ Ph	CN	p-CN	1
CH ₂ CH ₂ Ph	CN	p-NMe ₂	1
CH ₂ CH ₂ Ph	CN	p-NHMe	1
CH ₂ CH ₂ Ph	CN	p-CO ₂ H	1
CH ₂ CH ₂ Ph	CN	m-CO ₂ Et	1
CH ₂ CH ₂ Ph	CN	m-OMe	1
CH ₂ CH ₂ Ph	CN	p-NO ₂	1
CH ₂ CH ₂ Ph	F	p-NMe ₂	1
CH ₂ CH ₂ Ph	F	p-NHMe	1
CH ₂ CH ₂ Ph	Cl	p-NH ₂	2
CH ₂ CH ₂ Ph	Cl	p-Et	2
CH ₂ CH ₂ Ph	Br	p-Pr	2
CH ₂ CH ₂ Ph	Br	p-CH ₂ OMe	2

Table 6




R ⁴	R ⁶	R ¹²	n
H	H	p-MeO	0
c-Pr	H	p-MeO	1
c-Hex	H	p-MeO	1
Me	Cl	p-MeO	1
Et	Br	p-MeO	1
n-Pr	H	m-MeO	1
i-Pr	H	o-MeO	1
n-Bu	H	p-Me	1
n-Pen	H	p-Et	1
n-Hex	F	m-Et	1
COMe	NO ₂	o-Et	1
CONHMe	NO ₂	p-Cl	2
CONMe ₂	NO ₂	p-F	2
CF ₃	H	p-Ph	3
CH ₂ Ph	CN	p-OH	3
CH ₂ CH ₂ Ph	CN	p-NO ₂	4
CH ₂ CH ₂ Ph	CN	p-CN	1
CH ₂ CH ₂ Ph	CN	p-NMe ₂	1
CH ₂ CH ₂ Ph	CN	p-NHMe	1
CH ₂ CH ₂ Ph	CN	p-CO ₂ H	1
CH ₂ CH ₂ Ph	CN	m-CO ₂ Et	1
CH ₂ CH ₂ Ph	CN	m-OMe	1
CH ₂ CH ₂ Ph	CN	p-NO ₂	1
CH ₂ CH ₂ Ph	F	p-NMe ₂	1
CH ₂ CH ₂ Ph	F	p-NHMe	1
CH ₂ CH ₂ Ph	Cl	p-NH ₂	2
CH ₂ CH ₂ Ph	Cl	p-Et	2
CH ₂ CH ₂ Ph	Br	p-Pr	2
CH ₂ CH ₂ Ph	Br	p-CH ₂ OMe	2

Table 7




R ⁴	R ⁶	R ¹²	n
H	H	p-MeO	0
c-Pr	H	p-MeO	1
c-Hex	H	p-MeO	1
Me	Cl	p-MeO	1
Et	Br	p-MeO	1
n-Pr	H	m-MeO	1
i-Pr	H	o-MeO	1
n-Bu	H	p-Me	1
n-Pen	H	p-Et	1
n-Hex	F	m-Et	1
COMe	NO ₂	o-Et	1
CONHMe	NO ₂	p-Cl	2
CONMe ₂	NO ₂	p-F	2
CF ₃	H	p-Ph	3
CH ₂ Ph	CN	p-OH	3
CH ₂ CH ₂ Ph	CN	p-NO ₂	4
CH ₂ CH ₂ Ph	CN	p-CN	1
CH ₂ CH ₂ Ph	CN	p-NMe ₂	1
CH ₂ CH ₂ Ph	CN	p-NHMe	1
CH ₂ CH ₂ Ph	CN	p-CO ₂ H	1
CH ₂ CH ₂ Ph	CN	m-CO ₂ Et	1
CH ₂ CH ₂ Ph	CN	m-OMe	1
CH ₂ CH ₂ Ph	CN	p-NO ₂	1
CH ₂ CH ₂ Ph	F	p-NMe ₂	1
CH ₂ CH ₂ Ph	F	p-NHMe	1
CH ₂ CH ₂ Ph	Cl	p-NH ₂	2
CH ₂ CH ₂ Ph	Cl	p-Et	2
CH ₂ CH ₂ Ph	Br	p-Pr	2
CH ₂ CH ₂ Ph	Br	p-CH ₂ OMe	2

Table 8




R ¹	R ²	R ⁴	R ⁶	n
H	H	H	H	0
Me	Me	c-Pr	H	1
Me	Me	c-Hex	H	1
Me	Me	Me	Cl	1
Me	Me	Et	Br	1
Me	Me	n-Pr	H	1
Me	Me	i-Pr	H	1
Me	Me	n-Bu	H	1
Me	Me	n-Pen	H	1
Me	Me	n-Hex	F	1
Me	Me	COMe	NO ₂	1
Me	Me	CONHMe	NO ₂	2
Me	Me	CONMe ₂	NO ₂	2
Me	Me	CF ₃	H	3
Me	Me	CH ₂ Ph	CN	3
Me	Me	CH ₂ CH ₂ Ph	CN	4
Me	Me	CH ₂ CH ₂ Ph	NO ₂	1
Me	Me	CH ₂ CH ₂ Ph	NO ₂	2
Me	Me	CH ₂ CH ₂ Ph	NO ₂	3
Et	Et	CH ₂ CH ₂ Ph	CN	1
n-Pr	n-Pr	CH ₂ CH ₂ Ph	CN	1
i-Pr	i-Pr	CH ₂ CH ₂ Ph	CN	1
n-Bu	n-Bu	CH ₂ CH ₂ Ph	CN	1
i-Bu	i-Bu	CH ₂ CH ₂ Ph	F	1
t-Bu	t-Bu	CH ₂ CH ₂ Ph	F	1
n-Pen	n-Pen	CH ₂ CH ₂ Ph	Cl	2
n-Hex	n-Hex	CH ₂ CH ₂ Ph	Cl	2
CF ₃	CF ₃	CH ₂ CH ₂ Ph	Br	2
CH ₂ OCH ₃	CH ₂ OCH ₃	CH ₂ CH ₂ Ph	Br	2

Table 9




R ¹	R ²	R ⁴	R ⁶	n
H	H	H	H	0
Me	Me	c-Pr	H	1
Me	Me	c-Hex	H	1
Me	Me	Me	Cl	1
Me	Me	Et	Br	1
Me	Me	n-Pr	H	1
Me	Me	i-Pr	H	1
Me	Me	n-Bu	H	1
Me	Me	n-Pen	H	1
Me	Me	n-Hex	F	1
Me	Me	COMe	NO ₂	1
Me	Me	CONHMe	NO ₂	2
Me	Me	CONMe ₂	NO ₂	2
Me	Me	CF ₃	H	3
Me	Me	CH ₂ Ph	CN	3
Me	Me	CH ₂ CH ₂ Ph	CN	4
Me	Me	CH ₂ CH ₂ Ph	NO ₂	1
Me	Me	CH ₂ CH ₂ Ph	NO ₂	2
Me	Me	CH ₂ CH ₂ Ph	NO ₂	3
Et	Et	CH ₂ CH ₂ Ph	CN	1
n-Pr	n-Pr	CH ₂ CH ₂ Ph	CN	1
i-Pr	i-Pr	CH ₂ CH ₂ Ph	CN	1
n-Bu	n-Bu	CH ₂ CH ₂ Ph	CN	1
i-Bu	i-Bu	CH ₂ CH ₂ Ph	F	1
t-Bu	t-Bu	CH ₂ CH ₂ Ph	F	1
n-Pen	n-Pen	CH ₂ CH ₂ Ph	Cl	2
n-Hex	n-Hex	CH ₂ CH ₂ Ph	Cl	2
CF ₃	CF ₃	CH ₂ CH ₂ Ph	Br	2
CH ₂ OCH ₃	CH ₂ OCH ₃	CH ₂ CH ₂ Ph	Br	2

Table 10




R^4	R^6	R^{12}	$R^{12'}$	n
H	H	Br	Br	0
c-Pr	H	Br	Br	1
c-Hex	H	Br	Br	1
Me	Cl	Br	MeO	1
Et	Br	Br	Br	1
n-Pr	H	Br	Br	1
i-Pr	H	Br	MeO	1
n-Bu	H	Br	MeO	1
n-Pen	H	Br	MeO	1
n-Hex	F	Br	MeO	1
COMe	NO ₂	Br	MeO	1
CONHMe	NO ₂	Br	MeO	2
CONMe ₂	NO ₂	Br	Br	2
CF ₃	H	Br	Br	3
CH ₂ Ph	CN	Br	Br	3
CH ₂ CH ₂ Ph	CN	Br	Br	4
CH ₂ CH ₂ Ph	NO ₂	Br	Br	1
CH ₂ CH ₂ Ph	CN	Br	MeO	1
CH ₂ CH ₂ Ph	NO ₂	Br	MeO	1
CH ₂ CH ₂ Ph	CN	Cl	Cl	1
CH ₂ CH ₂ Ph	NO ₂	Cl	Cl	1
CH ₂ CH ₂ Ph	NO ₂	Cl	Cl	2
CH ₂ CH ₂ Ph	CN	Cl	MeO	1
CH ₂ CH ₂ Ph	F	Cl	MeO	1
CH ₂ CH ₂ Ph	F	Cl	MeO	2
CH ₂ CH ₂ Ph	Cl	Cl	MeO	1
CH ₂ CH ₂ Ph	Cl	Cl	MeO	2
CH ₂ CH ₂ Ph	Br	Cl	MeO	1
CH ₂ CH ₂ Ph	Br	Cl	MeO	2

Table 11



R ¹	R ²	R ⁹	R ⁶	n
H	H	p-MeO	H	0
Me	Me	p-MeO	H	1
Me	Me	p-MeO	H	2
Me	Me	p-MeO	H	3
Me	Me	H	H	1
Me	Me	m-MeO	H	0
Me	Me	o-MeO	H	1
Me	Me	p-Me	H	2
Me	Me	p-Et	H	3
Me	Me	m-Et	H	4
Me	Me	o-Et	H	2
Me	Me	p-Cl	H	2
Me	Me	p-F	H	2
Me	Me	p-Ph	H	2
Me	Me	p-OH	NO ₂	2
Me	Me	p-NO ₂	NO ₂	1
Me	Me	p-CN	NO ₂	2
Me	Me	p-NMe ₂	NO ₂	3
Me	Me	p-NHMe	NO ₂	4
Et	Et	p-CO ₂ H	NO ₂	2
n-Pr	n-Pr	m-CO ₂ Et	NO ₂	2
i-Pr	i-Pr	m-OMe	NO ₂	2
n-Bu	n-Bu	p-NO ₂	NO ₂	2
i-Bu	i-Bu	p-NMe ₂	NO ₂	2
t-Bu	t-Bu	p-NHMe	NO ₂	3
n-Pen	n-Pen	p-NH ₂	NO ₂	3
n-Hex	n-Hex	p-Et	NO ₂	3
CF ₃	CF ₃	p-Pr	NO ₂	3
CH ₂ OCH ₃	CH ₂ OCH ₃	p-CH ₂ OMe	NO ₂	3

Table 12



R ⁴	R ⁶	R ¹²	n
H	H	p-MeO	1
Me	H	p-MeO	1
Et	NO ₂	p-MeO	1
n-Pr	NO ₂	p-MeO	1
Et	H	p-MeO	4
i-Pr	H	m-MeO	0
i-Pr	H	o-MeO	1
CH ₂ CH ₂ Ph	H	p-MeO	1
n-Pen	H	p-Et	3
n-Hex	H	m-Et	4
COMe	H	o-Et	2
CONHMe	H	p-Cl	2
CONMe ₂	H	p-F	2
CF ₃	H	p-Ph	2
CH ₂ Ph	NO ₂	p-OH	2
CH ₂ CH ₂ Ph	NO ₂	p-MeO	1
Ph	NO ₂	p-MeO	1
CH ₂ Ph	NO ₂	p-MeO	1
H	NO ₂	p-MeO	1
Me	NO ₂	p-MeO	1
CH ₂ CH ₂ Ph	NO ₂	m-CO ₂ Et	1
CH ₂ CH ₂ Ph	NO ₂	m-OMe	1
CH ₂ CH ₂ Ph	NO ₂	p-NO ₂	1
CH ₂ CH ₂ Ph	NO ₂	p-NMe ₂	1
CH ₂ CH ₂ Ph	NO ₂	p-NHMe	1
CH ₂ CH ₂ Ph	NO ₂	p-NH ₂	1
CH ₂ CH ₂ Ph	NO ₂	p-Et	1
CH ₂ CH ₂ Ph	NO ₂	p-Pr	1
CH ₂ CH ₂ Ph	NO ₂	p-CH ₂ OMe	1

The compound according to the present invention has asymmetric carbon atoms at 3-position and 4-position, thus optical isomers thereof based on the asymmetric carbon atoms are present, which can be used in the application of the present invention similar to racemate thereof. Further, a cis or trans isomer based on configuration at 3-position and 4-position may be included, but the trans isomer is preferable.

Further, when the compounds can form their salts, the pharmaceutically acceptable salts can be also used as active ingredients.


As pharmaceutically acceptable salts, there may be mentioned hydrochlorides, hydrobromides, sulfates, methanesulfonates, acetates, benzoates, tartrates, phosphates, lactates, maleates, fumarates, malates, gluconates and salicylates, etc.

Preferably, there may be mentioned hydrochlorides and methanesulfonates.

Then, the preparation method of the compound according to the present invention is illustrated.

Of the compounds of the general formula (I), those wherein R^3 represents a hydroxyl group, which are the compounds of formula (I-a), can be obtained by reacting a compound of the general formula (2) with a compound (3) in an inert solvent, as shown in the following reaction scheme.

The compound of the general formula (2) can be synthesized according to known methods (methods described in J.M. Evans et al., J. Med. Chem. 1984, 27, 1127, J. Med. Chem. 1986, 29, 2194, J.T. North et al., J. Org. Chem. 1995, 60, 3397, as well as Japanese Patent Application Laid-Open No. Sho 56-57785, Japanese Patent Application Laid-Open No. Sho 56-57786, Japanese Patent Application Laid-Open No. Sho 58-188880, Japanese Patent Application Laid-Open No. Hei 2-141, Japanese Patent Application Laid-Open No. Hei 10-87650 and Japanese Patent Application Laid-Open No. Hei 11-209366, etc.).



In this scheme, $R^1, R^2, R^4, R^5, R^6, X$ and n are as defined above.

As the solvents used in the reaction of the compound of the general formula (2) with the compound (3), the following may be mentioned.

There may be mentioned sulfoxide type solvents exemplified by dimethylsulfoxide; amide type solvents exemplified by dimethylformamide or dimethylacetamide; ether type solvents exemplified by ethyl ether, dimethoxyethane or tetrahydrofuran; halogen type solvents exemplified by dichloromethane, chloroform and dichloroethane; nitrile type solvents exemplified by acetonitrile and propionitrile; aromatic hydrocarbon type solvents exemplified by benzene and toluene; hydrocarbon type solvents exemplified by hexane and heptane; and ester type solvents exemplified by ethyl acetate. Further, the reaction can be carried out in the absence of a solvent. Preferably, ether type solvents and nitrile type solvents may be mentioned.

The reaction temperature is generally from -80°C to the reflux temperature of the reaction solvent, preferably from -10°C to 100°C .

The molar ratio of the reaction materials is within the range of 0.5-20.0, preferably 1.0-10.0, for the compound (3)/the compound (2).

An acid catalyst may be used in the reaction.

As the acid catalysts used, there may be mentioned inorganic acids exemplified by hydrochloric acid and sulfuric acid, as well

as Lewis acids exemplified by aluminum chloride, titanium tetrachloride, boron trifluoride diethyl ether complex, perchloric acid, lithium perchlorate and ytterbium trifluoromethanesulfonate, etc.

Preferably, there may be mentioned sulfuric acid and perchloric acid.

Of the compounds of the general formula (I), those other than the compounds of formula (I-a) described above (those of the formula (I) wherein R^3 represents a C_{1-6} alkylcarbonyloxy group) can be prepared by the preparation methods similar to those described in Japanese Patent Application Laid-Open No. Sho 52-91866 and Japanese Patent Application Laid-Open No. Hei 10-87650, etc.

Syntheses of optically active compounds included in the compounds of the general formula (I) can be attained by utilizing optical resolution methods (Japanese Patent Application Laid-Open No. Hei 3-141286, U.S. Patent No. 5097037 and European Patent No. 409165). Further, syntheses of optically active compounds of the general formula (2) can be attained by utilizing asymmetrical synthetic methods (Japanese National Publication No. Hei 5-507645, Japanese Patent Application Laid-Open No. Hei 5-301878, Japanese Patent Application Laid-Open No. Hei 7-285983, European Patent Application Laid-open No. 535377, and U.S. Patent No. 5420314).

As described above, we, inventors, found that the compound of the general formula (I) has the strong prolongation effect on the functional refractory period. The prolongation effect on the functional refractory period is one of the functions of antiarrhythmic action and an important indicator that can be extrapolated to efficiency for clinical arrhythmia. Conventional antiarrhythmic agents having the prolongation effect on the functional refractory period as the main function (such as d-sotalol belonging to Class III of the antiarrhythmic agent classification according to Vaughan Williams) have quite dangerous arrhythmic inducing actions that can result in sudden death such as torsades de pointes based on extension of ventricular muscle action potential relating to the prolongation effect on the functional refractory period, which become the therapeutic problems for arrhythmia based

on atrium (such as supraventricular tachycardia, atrial flutter and atrial fibrillation). In order to solve the problems, we, inventors, carried out searching and studying of compounds having the prolongation effect on the functional refractory period more selective for atrium muscle than for ventricular muscle, and found that the compound of the general formula (I) has the prolongation effect on the functional refractory period selective for atrium muscle without any influence on the functional refractory period of ventricular muscle and action potential parameters. The difference between the present invention by the inventors and the known techniques is to provide the prolongation effect on the functional refractory period selective for atrium muscle by the compound, which is shown by the following facts; without any influence on the action potential sustaining period of removed ventricular muscle and without any influence on the electrocardiogram QT of anesthetized animal. From the above, the compounds of the present invention have no arrhythmic inducing action in ventricular muscle, thus they can provide possibilities of more safe uses for arrhythmia based on atrium muscle than known techniques. The technique according to the present invention is useful for therapeutic or preventive uses as anti-atrial fibrillation agents, anti-atrial flutter agents and anti-atrial tachycardia agents relating to paroxysmal, chronic, preoperative, intraoperative or postoperative atrial arrhythmia, prevention of proceeding to embolus based on atrial arrhythmia, prevention of proceeding to ventricular arrhythmia or tachycardia originated from atrial arrhythmia or tachycardia, and prevention of the life prognosis worsening based on the preventive action for atrial arrhythmia or tachycardia which can be proceeded to ventricular arrhythmia or tachycardia.

The present invention provides a pharmaceutical composition or veterinary pharmaceutical composition containing the compound of the generally formula (I) in an effective amount for these treatments.

As administering forms of the compound according to the present invention, there may be mentioned parenteral

administrations by means of injections (subcutaneous, intravenous, intramuscular and intraperitoneal injections), ointments, suppositories and aerosol, or oral administrations by means of tablets, capsules, granules, pills, syrups, solutions, emulsions and suspensions, etc.

The above-mentioned pharmaceutical or veterinary pharmaceutical composition contains the compound according to the present invention in an amount of about 0.01-99.5%, preferably about 0.1-30%, of the total composition weight.

In addition to the compound according to the present invention or the composition containing the compound, other pharmaceutically or veterinary pharmaceutically active compounds may be contained.

Further, these compositions may contain the plurality of compounds according to the present invention.

A clinical administration amount varies depending on age, weight and sensitivity of the patient, extent of condition of the patient, etc. and an effective administration amount is generally about 0.003-1.5 g, preferably 0.01-0.6 g, per day for adult. If necessary, however, the amount outside of the above-mentioned range may be used.

The compound according to the present invention is formulated for administration by conventional pharmaceutical means.

That is, tablets, capsules, granules and pills for oral administration are prepared by using excipients such as sucrose, lactose, glucose, starch and mannitol; binders such as hydroxypropyl cellulose, syrup, gum arabic, gelatin, sorbitol, tragacanth, methyl cellulose and polyvinyl pyrrolidone; disintegrators such as starch, carboxymethyl cellulose or its calcium salt, crystal cellulose powder and polyethylene glycol; lubricants such as talc, magnesium or calcium stearate, and silica; lubricating agents such as sodium laurate and glycerol, etc.

Injections, solutions, emulsions, suspensions, syrups and aerosols are prepared by using solvents for the active ingredients such as water, ethyl alcohol, isopropyl alcohol, propylene glycol, 1,3-butylene glycol and polyethylene glycol; surfactants such as sorbitan fatty acid ester, polyoxyethylene sorbitan fatty acid

ester, polyoxyethylene fatty acid ester, polyoxyethylene ether of hydrogenated castor oil and lecithin; suspending agents such as carboxymethyl sodium salt, cellulose derivatives such as methyl cellulose, tragacanth, and natural rubbers such as gum arabic; and preserves such as p-hydroxybenzoic acid esters, benzalkonium chloride and sorbic acid salts, etc.

For ointments that are transdermally adsorptive pharmaceuticals, white vaseline, liquid paraffin, higher alcohols, Macrogol ointments, hydrophilic ointments and aqueous gel-type bases are, for example, used.

Suppositories are prepared by using, for example, cocoa fats, polyethylene glycol, lanolin, fatty acid triglyceride, coconut oil and Polysorbate etc.


Best Mode for Carrying Out the Invention

The present invention is illustrated in detail by the Examples as follows, but the present invention is not limited to these Examples.

[Synthesis Examples]

Synthesis example 1

(3R*, 4S*)-3,4-dihydro-2,2-dimethyl-6-(4-methoxyphenylacetyl-amino)-7-nitro-2H-1-benzopyran-3,4-diol



To a mixed solution of (3R*, 4R*) -3,4-epoxy-3,4-dihydro-2,2-dimethyl-6-(4-methoxyphenylacetyl-amino)-7-nitro-2H-1-benzopyran (the synthetic method of which is described in Japanese Patent Laid-Open Publication No. Hei 10-87650) (300 mg, 0.78 mmol, 99%ee, $[\alpha]_D^{25} +6.2$ (C 0.70, CHCl₃)) in acetonitrile (2 mL) and a 70% aqueous perchloric acid solution (1 mL), water (1 mL) was added at the room temperature and stirred at the room temperature for

2 hours. After adding a 10% aqueous sodium carbonate solution to the solution, an aqueous saturated sodium chloride solution was added. Then the resulting solution was extracted with ethyl acetate, and dried over anhydrous sodium sulfate. After the solvent was distilled off, the residue was recrystallized (from methanol-water), to obtain the intended substance (yield: 81%).


$^1\text{H-NMR}$ (CDCl_3) δ : 1.27 (s, 3H), 1.46 (s, 3H), 3.60-3.65 (m, 1H), 3.73 (s, 2H), 3.79 (s, 3H), 3.90-3.94 (m, 2H), 4.60-4.62 (m, 1H), 6.95 (d, $J = 8.6$ Hz, 2H), 7.26 (d, $J = 8.6$ Hz, 2H), 7.57 (s, 1H), 8.79 (s, 1H), 9.93 (s, 1H).

MS (EI) m/z ; 402 $[\text{M}]^+$, 356, 254, 182, 148, 122 (bp).

The following compounds were obtained in the similar manner.

Synthesis example 2

(3R*, 4S*)-3,4-dihydro-2,2-dimethyl-4-methoxy-6-(4-methoxyphenylacetyl-amino)-7-nitro-2H-1-benzopyran-3-ol




Yield: 39 %

$^1\text{H-NMR}$ (CDCl_3) δ : 1.27 (s, 3H), 1.46 (s, 3H), 2.34 (m, 1H), 3.64 (s, 3H), 3.73 (s, 2H), 3.83 (s, 3H), 3.80-3.85 (m, 1H), 4.31-4.34 (m, 1H), 6.95 (d, $J = 8.8$ Hz, 2H), 7.26 (d, $J = 8.8$ Hz, 2H), 7.58 (s, 1H), 8.75 (s, 1H), 9.9 (s, 1H).

MS (EI) m/z ; 416 $[\text{M}]^+$, 268, 148, 121 (bp). ; mp. 104 °C.

Synthesis example 3

(3R*, 4S*)-4-ethoxy-3,4-dihydro-2,2-dimethyl-6-(4-methoxyphenylacetyl-amino)-7-nitro-2H-1-benzopyran-3-ol



Yield: 53 %


Orange amorphous substance

$^1\text{H-NMR}$ (CDCl_3) δ : 1.27 (s, 3H), 1.35 (t, $J = 7.0$ Hz, 3H), 1.46 (s, 3H), 2.33 (m, 1H), 3.73 (s, 2H), 3.83 (s, 3H), 3.76-4.0 (m, 3H), 4.36 (d, $J = 7.9$ Hz, 1H), 6.96 (d, $J = 8.6$ Hz, 2H), 7.26 (d, $J = 8.6$ Hz, 2H), 7.57 (s, 1H), 8.79 (s, 1H), 9.93 (s, 1H).

MS (EI) m/z ; 430 $[\text{M}]^+$, 282, 148, 121 (bp).

Synthesis example 4

(3R*, 4S*)-3,4-dihydro-2,2-dimethyl-4-propoxy-6-(4-methoxyphenylacetamino)-7-nitro-2H-1-benzopyran-3-ol



Yield: 40 %


mp. 115-116 °C.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.01 (t, $J = 7.4$ Hz, 3H), 1.27 (s, 3H), 1.45 (s, 3H), 1.73 (q, $J = 6.6$ Hz, 2H), 2.35 (m, 1H), 3.73 (s, 2H), 3.68-3.85 (m, 4H), 3.83 (s, 3H), 6.95 (d, $J = 8.6$ Hz, 2H), 7.26 (d, $J = 8.6$ Hz, 2H), 7.57 (s, 1H), 8.78 (s, 1H), 9.92 (s, 1H).

MS (EI) m/z ; 444 $[\text{M}]^+$, 296, 148, 121 (bp).

Synthesis example 5

Trans-3,4-dihydro-2,2-dimethyl-6-(4-methoxyphenylacetamino)-4-(2-phenylethoxy)-2H-1-benzopyran-3-ol



To a solution of 3,4-epoxy-3,4-dihydro-2,2-dimethyl-6-(4-methoxyphenylacetylamino)-2H-1-benzopyran (200 mg, 0.68 mmol) and phenethyl alcohol (0.32 mL, 2.72 mmol) in acetonitrile (1.0 mL), sulfuric acid (0.01 mL) was added and stirred at the room temperature for 2 hours. Thereto, ice-cooled ethyl acetate was added and the formed crystals were filtered, to obtain the intended substance as white solid (yield: 77%).

m.p. : 161.0-162.0 °C


$^1\text{H-NMR}$ (CDCl_3) δ : 1.17 (s, 3H), 1.35 (s, 3H), 2.93 (t, 2H, $J = 6.4$ Hz), 3.64 (d, 1H, $J = 7.6$ Hz), 3.66 (s, 2H), 3.83 (s, 3H), 3.84 - 3.90 (m, 1H), 3.94 - 3.99 (m, 1H), 4.29 (d, 1H, $J = 7.6$ Hz), 6.68 (d, 2H, $J = 8.8$ Hz), 6.84 (s, 1H), 6.94 (d, 2H, $J = 8.8$ Hz), 7.11 (d, 1H, $J = 2.4$ Hz), 7.19 - 7.32 (m, 6H).

MS (EI) m/z ; 461 $[M]^+$, 389, 340, 192 (bp).

The following compounds were obtained in the similar manner.

Synthesis example 6

Trans-3,4-dihydro-2,2-dimethyl-6-(4-methoxyphenyl)acetylamino-7-nitro-4-(2-phenylethoxy)-2H-1-benzopyran-3-ol



mp. : 125.2-125.5 °C


$^1\text{H-NMR}$ (CDCl_3) δ : 1.17 (s, 3H), 1.36 (s, 3H), 1.84 (d, $J = 4.0$ Hz, 1H), 2.98-3.01 (m, 2H), 3.60 (dd, $J = 4.0, 8.2$ Hz, 1H), 3.74 (s,

2H), 3.83 (s, 3H), 3.90-3.96 (m, 1H), 4.15-4.21 (m, 1H) 4.32 (dd, $J = 0.9, 8.2$ Hz, 1H), 6.94-7.55 (m, 10H), 8.76 (d, $J = 0.9$ Hz, 1H), 9.89 (bs, 1H)

MS (EI) m/z ; 507 $[m]^+$, (bp), 358.

Synthesis example 7

Trans-3,4-dihydro-2,2-dimethyl-6-(4-methoxyphenyl)acetylamino-7-nitro-4-phenyloxy-2H-1-benzopyran-3-ol




mp. : 178.0-178.6 °C

$^1\text{H-NMR}$ (DMSO- d_6) δ : 1.31 (s, 3H), 1.37 (s, 3H), 3.52 (s, 2H), 3.72 (s, 3H), 3.77 (dd, $J = 5.7, 6.0$ Hz, 1H), 5.31 (d, $J = 6.0$ Hz, 1H), 5.92 (d, $J = 5.7$ Hz, 1H), 6.86-7.48 (m, 11H), 10.16 (bs, 1H)

MS (EI) m/z ; 478 $[M]^+$, 237 (bp).

Synthesis example 8

Trans-3,4-dihydro-2,2-dimethyl-6-(4-methoxyphenyl)acetylamino-7-nitro-4-benzyloxy-2H-1-benzopyran-3-ol



mp. : 117.8-119.1 °C


$^1\text{H-NMR}$ (DMSO- d_6) δ : 1.24 (s, 3H), 1.38 (s, 3H), 3.56 (s, 2H), 3.74 (s, 3H), 3.77-3.80 (m, 1H), 4.48 (d, $J = 7.0$ Hz, 1H), 4.80 (d, $J = 11.4$ Hz, 1H), 4.90 (d, $J = 11.4$ Hz, 1H), 5.78 (d, $J = 5.7$ Hz,

1H), 6.88-7.58 (m, 11H), 10.15 (bs, 1H)

MS (EI) m/z; 492 [M]⁺, 123 (bp).

Synthesis example 9

Trans-3,4-dihydro-2,2-dimethyl-7-nitro-4-(2-phenylethoxy)-6-(4-methoxyurea)-2H-1-benzopyran-3-ol



¹H-NMR (CDCl₃) δ : 1.18 (s, 3H), 1.36 (s, 3H), 2.95-2.98 (m, 2H), 3.59-3.60 (m, 1H), 3.89-3.95 (m, 1H), 4.13 (d, J = 8.4 Hz, 1H) 4.29 (d, J = 8.4 Hz, 1H), 7.13-7.54 (m, 12H), 8.53 (s, 1H), 9.50 (s, 1H)

MS (EI) m/z; 477 [M]⁺, (bp), 120.

[Preparation Examples]

Preparation Example 1

Tablet:

a compound according to the invention	10 g
lactose	260 g
crystal cellulose powder	600 g
corn starch	350 g
hydroxypropyl cellulose	100 g
CMC-Ca	150 g
<u>magnesium stearate</u>	<u>30 g</u>
total	1,500 g

The above-mentioned compounds were mixed by a usual method and thereafter 10,000 sugar-coated tablets each containing 1 mg of the active ingredient per a tablet were prepared.

Preparation Example 2

Capsule:

a compound according to the invention	10 g
lactose	440 g
crystal cellulose powder	1,000 g
<u>magnesium stearate</u>	<u>50 g</u>
total	1,500 g

The above-mentioned compounds were mixed by a usual method and thereafter filled in gelatin capsules, to prepare 10,000 capsules each containing 1 mg of the active ingredient per a capsule.

Preparation Example 3

Soft capsule:

a compound according to the invention	10 g
PEG 400	479 g
saturated fatty acid triglyceride	1,500 g
peppermint oil	1 g
<u>Polysorbate 80</u>	<u>10 g</u>
Total	2,000 g

The above-mentioned compounds were mixed by a usual method and thereafter filled in No.3 soft gelatin capsules, to prepare 10,000 soft capsules each containing 1 mg of the active ingredient per a capsule.

Preparation Example 4

Ointment:

a compound according to the invention	1.0 g
liquid paraffin	10.0 g
cetanol	20.0 g
white vaseline	68.4 g
ethylparaben	0.1 g
<u>1-menthol</u>	<u>0.5 g</u>
total	100.0 g

The above-mentioned compounds were mixed by a usual method to obtain 1% ointment.

Preparation Example 5

Suppository:

a compound according to the invention	1 g
Witepsol H15*	478 g
Witepsol W35*	520 g
<u>Polysorbate 80</u>	<u>1 g</u>
Total	1,000 g

(* trade name Witepsol for triglyceride type compounds)

The above-mentioned compounds were melt-mixed by a usual method, poured into suppository containers and cooled to solidify, thereby 1,000 suppositories (1 g) each containing 1 mg of the active ingredient per a suppository were prepared.

Preparation Example 6

Injection:

a compound according to the invention	1 mg
distilled water for injection	5 mL

It is used by dissolving when applied.

[Pharmacological Test Example]Effects of compound on the functional refractory period in guinea-pig left atrium muscle and right ventricular papillary muscleTest method

Hearts were removed from guinea-pigs, and left atrium muscle or right ventricular papillary muscle were isolated therefrom in a Krebs-Henseleit solution aerated with 95% O₂ + 5% CO₂. The samples were stimulated electrically at a rate of 1 Hz and a voltage of 1.5 times of the threshold value reacted to stimulation (basic stimulation; S1) by using an electric stimulating apparatus. The contraction occurred at that time was recorded by a thermal stylus recorder via a FD pickup and a strain pressure amplifier. The functional refractory period is defined as the shortest time interval between S1 resulting from determinable contraction and an extra stimulation (S2). The time interval between S1 and S2 in the left atrium muscle sample was started from 150 msec, decreased

in 10 msec steps until 100 msec, and thereafter 5 msec steps to the functional refractory period. For the right ventricular papillary muscle sample, it was started from 300 msec and decreased in 10 msec steps until the functional refractory period. Herein, S2 was set at twice of the threshold value which reacted to stimulation. The experimental temperature was $36 \pm 1^\circ\text{C}$. Herein, the solvent did not influence on any of the functional refractory periods for left atrium muscle and right ventricular papillary muscle. After determining the basic value before addition of the compound, the compound was added cumulatively, incubated for 15 minutes for respective concentration, and thereafter the functional refractory period was determined.

Results

Compounds according to the present invention exhibited strong prolongation effect on the functional refractory period (FRP) on atrium muscle.


Table 13

Synthesis example No.	Prolongation effect on FRP $\text{EC}_{20}(\mu\text{M})$
6	2.4
9	21.9

Compounds according to the present invention exhibit strong prolongation effect on the functional refractory period, thus they are useful for improvement of arrhythmia. Therefore, the present invention can provide useful antiarrhythmic agents.

CLAIMS

1. A 4-oxybenzopyran derivative of the formula (I)



wherein, R^1 and R^2 represent each independently a hydrogen atom; a C_{1-6} alkyl group in which said alkyl group may be optionally substituted with a halogen atom, a C_{1-6} alkoxy group or a hydroxyl group; or a phenyl group in which said phenyl group may be optionally substituted with a halogen atom, a hydroxyl group, a nitro group, a cyano group, a C_{1-6} alkyl group or a C_{1-6} alkoxy group;

R^3 represents a hydroxyl group or a C_{1-6} alkylcarbonyloxy group;


R^4 represents a hydrogen atom, a C_{3-6} cycloalkyl group, a C_{1-6} alkyl group, a C_{1-6} alkylcarbonyl group, a C_{1-6} alkylaminocarbonyl group or a di- C_{1-6} alkylaminocarbonyl group in which said C_{1-6} alkyl group, said C_{1-6} alkylcarbonyl group, said C_{1-6} alkylaminocarbonyl group and said di- C_{1-6} alkylaminocarbonyl group may be each optionally substituted with a halogen atom, a C_{1-6} alkoxy group, a C_{1-6} alkoxy group substituted by a halogen atom; a carboxyl group, a C_{1-6} alkoxycarbonyl group, a hydroxyl group, or an aryl group or a heteroaryl group, in which said aryl group and said heteroaryl group may be optionally substituted with $(R^9)_x$, in which R^9 represents a halogen atom, a hydroxyl group, a C_{1-6} alkyl group, a C_{1-6} alkyl group substituted by a halogen atom or a C_{1-6} alkoxy group; a C_{1-6} alkoxy group, a C_{1-6} alkoxy group substituted by a halogen atom; or R^9 represents a nitro group, a cyano group, a formyl group, a formamide group, an amino group, a C_{1-6} alkylamino group, a di- C_{1-6} alkylamino group, a C_{1-6} alkylcarbonylamino group, a C_{1-6} alkylsulfonylamino group, an aminocarbonyl group, a C_{1-6} alkylaminocarbonyl group, a di- C_{1-6} alkylaminocarbonyl group, a C_{1-6}

alkylcarbonyl group, a C_{1-6} alkoxy carbonyl group, an aminosulfonyl group, a C_{1-6} alkylsulfonyl group, a carboxyl group or an arylcarbonyl group, r represents an integer of 1-3 and each R^9 may same or different if r represents 2 or 3; or R^4 represents an aryl group or a heteroaryl group in which said aryl group and said heteroaryl group may be optionally substituted with $(R^{10})_q$ in which R^{10} has the same meaning as R^9 , q represents an integer of 1-3, and each R^{10} may be same or different if q represents 2 or 3;

n represents an integer of 0-4;

X represents $-C(=O)NR^7-$, $-NR^8-$, $-NHC(=O)NH-$ or $-S(O)_2NH-$ (in which R^7 and R^8 represent each independently a hydrogen atom or a C_{1-6} alkyl group);

R^5 represents a hydrogen atom; or a group of the formula:



in which R^{12} represents a hydrogen atom; a halogen atom; a C_{1-6} alkyl group in which the alkyl group may be optionally substituted with a halogen atom or a C_{1-6} alkoxy group; a C_{1-6} alkoxy group in which said alkoxy group may be optionally substituted with a halogen atom; a phenyl group in which said phenyl group may be optionally substituted with a halogen atom, a hydroxyl group, a C_{1-6} alkyl group

or a C₁₋₆ alkoxy group; or R¹² represents a hydroxyl group, a nitro group, a cyano group, a formyl group, a formamide group, an amino group, a C₁₋₆ alkylamino group, a di-C₁₋₆ alkylamino group, a C₁₋₆ alkylcarbonylamino group, a C₁₋₆ alkylsulfonylamino group, an aminocarbonyl group, a C₁₋₆ alkylaminocarbonyl group, a di-C₁₋₆ alkylaminocarbonyl group, a C₁₋₆ alkylcarbonyl group, a C₁₋₆ alkoxycarbonyl group, an aminosulfonyl group, a C₁₋₆ alkylsulfonyl group, a carboxyl group or an arylcarbonyl group, m represents an integer of 1-3, and each R¹² may be same or different if m represents 2 or 3, and R¹³ represents a hydrogen atom or a C₁₋₆ alkyl group; and


R⁶ represents a hydrogen atom, a halogen atom, a nitro group or a cyano group;
or a pharmaceutically acceptable salt thereof.

2. A 4-oxybenzopyran derivative or pharmaceutically acceptable salt thereof according to claim 1, wherein R¹ and R² represent both methyl groups and R³ represents a hydroxyl group.

3. A 4-oxybenzopyran derivative or pharmaceutically acceptable salt according to claim 2, wherein R⁶ represents a hydrogen atom or a nitro group.

4. A 4-oxybenzopyran derivative or pharmaceutically acceptable salt thereof according to claim 3, wherein X represents -C(=O)NH-, -C(=O)NMe-, -S(O)₂NH- or -NHC(=O)NH-.

5. A 4-oxybenzopyran derivative or pharmaceutically acceptable salt thereof according to claim 4, wherein X represents -C(=O)NH- and R⁵ represents



6. A 4-oxybenzopyran derivative or pharmaceutically acceptable salt thereof according to claim 5, wherein R⁴ represents a C₁₋₆ alkyl group.

7. A drug characterized by comprising a 4-oxybenzopyran derivative or pharmaceutically acceptable salt thereof according to claim 1 as an active ingredient.

8. A drug for treating arrhythmia characterized by comprising a 4-oxybenzopyran derivative or pharmaceutically acceptable salt thereof according to claim 1 as an active ingredient.

INTERNATIONAL SEARCH REPORT

Internat. Application No

PCT/JP 00/06497

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D311/22 C07D405/12 A61K31/353 A61P9/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 2 204 868 A (SANDOZ) 23 November 1988 (1988-11-23) page 39; claims; example 29	1,2,7,8
X	CHEMICAL ABSTRACTS, vol. 126, no. 17, 1997 Columbus, Ohio, US; abstract no. 225191h, SUN, HONG BIN: "FACILE SYNTHESIS OF TRANS-3,4-DIOL DERIVATIVES OF SUBSTITUTED CHROMANS" page 566; XP002160972 abstract & CHIN. CHEM. LETT., vol. 8, no. 1, 1997, pages 1-4, CHINA	1,2

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *G* document member of the same patent family

Date of the actual completion of the international search

21 February 2001

Date of mailing of the international search report

08/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Francois, J

INTERNATIONAL SEARCH REPORT

Internat. Application No

PCT/JP 00/06497

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 112 972 A (R.GERICKE) 12 May 1992 (1992-05-12) the whole document ----	1,2,7,8
A	EP 0 677 519 A (JAPAN TOBACCO) 18 October 1995 (1995-10-18) claims; examples 17,24 -----	1,7,8

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internat. Application No

PCT/JP 00/06497

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 2204868 A	23-11-1988	AT 125188 A	15-06-1993
		AU 614285 B	29-08-1991
		AU 1615888 A	17-11-1988
		BE 1002436 A	12-02-1991
		CA 1312866 A	19-01-1993
		CH 674984 A	15-08-1990
		DE 3815325 A	24-11-1988
		DK 265188 A	17-11-1988
		ES 2012527 A	01-04-1990
		FI 882248 A	17-11-1988
		FR 2615191 A	18-11-1988
		HU 50151 A, B	28-12-1989
		IT 1219940 B	24-05-1990
		JP 63303977 A	12-12-1988
		NL 8801258 A	16-12-1988
		NZ 224604 A	26-04-1991
		PT 87492 A, B	31-05-1989
		SE 467924 B	05-10-1992
		SE 8801807 A	17-11-1988
		US 5071871 A	10-12-1991
		ZA 8803450 A	31-01-1990
US 5112972 A	12-05-1992	DE 3918041 A	06-12-1990
		AT 110072 T	15-09-1994
		AU 623780 B	21-05-1992
		AU 5600790 A	06-12-1990
		CA 2017961 A	02-12-1990
		DE 59006817 D	22-09-1994
		DK 400430 T	12-09-1994
		EP 0400430 A	05-12-1990
		ES 2057262 T	16-10-1994
		FI 95701 B	30-11-1995
		HU 58323 A	28-02-1992
		IE 901988 A	02-01-1991
		IL 94256 A	12-04-1994
		JP 3020275 A	29-01-1991
		NO 175976 B	03-10-1994
		NZ 233912 A	28-10-1992
		PT 94225 A, B	08-02-1991
		ZA 9004227 A	27-03-1991
EP 677519 A	18-10-1995	KR 170534 B	18-02-1999
		US 5719155 A	17-02-1998
		CA 2153498 A	18-05-1995
		WO 9513272 A	18-05-1995
		JP 2649610 B	03-09-1997